

Preventive Services and Coding

Triple S Salud covers the following preventive services in Puerto Rico with no cost sharing and as required by the ACA (Affordable Care Act) and the Puerto Rico Health Insurance Code for non-grandfathered plans. Preventive services are those provided while the person does not present symptoms of a medical condition.

If a service does not meet all the coding criteria, a deductible, copayment, or coinsurance may apply, as described in the policy. Please review your policy or consult your plan's benefits administrator to determine if you need to pay for any of these services.

The following table is a reference to inform you about the preventive services you may have access to. However, it is not a guarantee of payment since these services may change due to legal or regulatory requirements. It is important that the provider complies with standard coding guidelines when billing the service to be considered preventive. If you have any questions, you can contact Customer Services, your plan administrator, or the Provider Services department if you are a physician or healthcare provider.

Preventives Care for Adults

Preventive Services for Adults	Procedure Codes and Facility Codes	Diagnosis Codes
Abdominal Aortic Aneurysm One (1) service per ultrasonogram for abdominal aortic aneurysm (AAA) screening, for members 65 to 75 years of age who currently are or were smokers at some point	Procedure: 76706 Facility: 0402	Z13.6 Z87.891
Screening for Anxiety Disorders The USPSTF recommends screening for anxiety disorders in adults (64 years of age or younger), including pregnant or postpartum individuals.	99385 99386	Z13.31 Z13.32 Z13.39
Colorectal cancer According to administrative order #334 of the Department of Health, one annual occult blood test for colorectal cancer screening is covered for adults 40 years of age and older. If the person has a family history of colorectal cancer, the annual screening is performed via colonoscopy instead of the occult blood test. The USPSTF recommends performing the colorectal cancer screening via occult blood test, sigmoidoscopy, colonoscopy, or serological test in adults 45 to 75 years old. The risks and benefits of these screening methods vary. The USPSTF also recommends performing a follow-up colonoscopy after a positive result on a non-invasive test. This is a screening test, and patients will not incur any out-of-pocket expenses.	Procedure: 82274 82270 Facility: 0301 Procedure: G0104 G0105 G0121 G0328 Facility: 0369	Z12.11, Z12.12 Z12.11, Z12.12
Depression and Suicide Risk Screening for Adults The USPSTF recommends screening for depression in the adult population, including individuals during pregnancy or postpartum, as well as older adults (65 years of age or older).	99385 99386 99387 99395 99396 99397 96160 96161	Z13.31 Z13.32 Z13.39
Fall prevention in senior citizens: exercise or physical therapy Exercises and physical therapy to prevent falls in adults over 65 years old who are at risk of falling.	99401 99402 99403 99404	Z91.81

Preventive Services for Adults	Procedure Codes and Facility Codes	Diagnosis Codes
<p>Healthy diet and physical activity as a form of prevention for cardiovascular disease in adults at cardiovascular risk. Offering and referring adults who are overweight or obese and who have additional risk factors for cardiovascular disease to intensive behavioral counseling interventions, to promote a healthy diet and physical activity to prevent cardiovascular diseases.</p>	97802 97803 97804 99401 99402 99403	Z71.3
<p>Hepatitis B virus screening The USPSTF recommends screening for the Hepatitis B virus in adults at high risk for infection.</p>	Procedure 87340 Facility 0306	Z11.59
<p>Hepatitis C virus screening: teenagers and adults Screening for hepatitis C (HCV) infections in adults from 18 to 79 years old.</p>	Procedure 86803 Facility 0306	Z11.59
<p>Hypertension screening for members who have not been diagnosed with the condition Hypertension screening for adults 18 years of age and older. Readings should be obtained outside the clinical setting to confirm the diagnosis before starting treatment.</p>	99385 99386 99387 99395 99396 99397	Z13.6
<p>Prevention of HIV Acquisition: Pre-exposure Prophylaxis: adolescents and adults at increased risk for HIV The USPSTF recommends that physicians prescribe pre-exposure prophylaxis using antiretroviral therapy to individuals who are at increased risk of contracting HIV in order to decrease the risk of acquiring HIV. See the USPSTF Practice Considerations section for more information on identifying persons at increased risk and effective antiretroviral therapy.</p>	N/A	N/A
<p>Human Immunodeficiency Virus (HIV) screening test: teenagers and adults that are not currently pregnant Human Immunodeficiency Virus (HIV) screening for adults 13 to 65 years old, as well as younger teenagers and older adults with high risk. As required by Act No. 45-2016, this includes one HIV test per year as part of the routine medical evaluation, except for pregnant members to whom USPSTF requirements apply.</p>	Procedure 86701 86702 86703 86689 87390 Facility 0302	Z11.4, Z11.59, Z70.0, Z70.1, Z70.2, Z70.3, Z70.8, Z70.9, Z71.7, Z72.51, Z72.52, Z72.53
<p>Immunization Vaccines. The recommended dosages, ages, and population vary: Hepatitis A, Hepatitis B, Herpes Zoster, Human Papilloma Virus, Influenza, MMR, Meningococcus, Pneumococcus, Tetanus, Diphtheria, Whooping Cough, and Chickenpox, Haemophilus influenza type B. Catch-up vaccines are covered. COVID-19 vaccine for persons 19 years of age and older adults, as recommended by the Advisory Committee on Immunization Practices (ACIP) of the Centers for Disease Control and Prevention (CDC) and in accordance with section 2.050 (c) of the Puerto Rico Health Insurance Code.</p>	90380	Z29.11
	90381	Z29.11
	90620	D57.1, D57.20, D57.21, D57.212, D57.40, D57.411, D57.412, D57.80, D57.811, D57.812, D84.1, Q89.01, Z20.811, Z23, Z90.81
	90621	D57.1, D57.20, D57.21, D57.212, D57.40, D57.411, D57.412, D57.80, D57.811,

Preventive Services for Adults	Procedure Codes and Facility Codes	Diagnosis Codes
<p><i>The Mpox vaccine is recommended for anyone at risk of Mpox infection.</i></p>		D57.812, D84.1,Q89.01 Z20.811, Z23, Z90.82
	90630	Z23
<p>The vaccine against respiratory syncytial virus (RSV) is recommended using seasonal administration for pregnant individuals and older adults over 60 years of age. See CDC recommendations.</p>	90651	Z23
	90653	Z23
	90671	Z23
	90677	Z23
	90678	Z29.11
	90747	Z23
	90750	Z23
	91304	Z23
	91318	Z23
	91319	Z23
	91320	Z23
	91321	Z23
	91322	Z23
	90632	Z23
	90633	Z23
	90634	Z23
	90636	Z23
	90647	Z23
	90648	Z23
	90649	Z23
	90650	Z23
	90654	Z23
	90655	Z23
	90656	Z23
	90657	Z23
	90658	Z23
	90661	Z23
	90662	Z23
	90670	Z23
	90673	Z23
	90674	Z23
	90680	Z23
	90681	Z23
90682	Z23	
90685	Z23	
90686	Z23	
90687	Z23	
90688	Z23	
90696	Z23	
90698	Z23	
90700	Z23	
90702	Z23	
90707	Z23	
90713	Z23	
90714	Z23	
90715	Z23	
90716	Z23	

Preventive Services for Adults	Procedure Codes and Facility Codes	Diagnosis Codes
	90723	Z23
	90732	Z23
	90733	Z23
	90734	Z23
	90740	Z23
	90743	Z23
	90744	Z23
	90746	Z23
	90748	Z23
	90671	Z29.11
	90678	Z29.11
<p>Detection of latent tuberculosis infection The USPSTF recommends screening tests for latent tuberculosis infection in high-risk populations. See the USPSTF "Risk Assessment" section for additional information on adults at increased risk.</p>	86580	Z11.1
<p>Lung cancer screening Annual lung cancer screening through computerized tomography for adults from 50 to 80 years old with a history of smoking twenty (20) packs per year, who are currently smoking or stopped smoking within the last 15 years. The screening will be discontinued once the person has stopped smoking for 15 consecutive years or develops a health problem that substantially limits their life expectancy or the likelihood of undergoing lung surgery to cure the disease.</p>	71250 71271	Z12.2, Z87.891
<p>Obesity screening and counseling for adults Physicians may offer or refer patients with a Body Mass Index (BMI) of 30 kg/m² or more to intensive multi-component behavioral interventions.</p>	99385 99386 99387 99395 99396 99397	E66.01, Z68.41, Z68.42, Z68.43, Z68.44, Z68.45
<p>Screening for prediabetes and type 2 diabetes in asymptomatic adults aged 35 to 70 years old and who are overweight or obese The USPSTF recommends screenings for prediabetes and type 2 diabetes in adults aged 35 to 70 years old who are overweight or obese. Physicians must offer or refer patients with prediabetes to effective preventive interventions.</p>	82947 82950 83036	E66.01, E66.09, E66.1, E66.2, E66.3, E66.8, E66.9, Z13.1, Z68.25, Z68.26, Z68.27, Z68.28, Z68.29, Z68.30, Z68.31, Z68.32, Z68.33, Z68.34, Z68.35, Z68.36, Z68.37, Z68.39, Z68.41, Z68.42, Z68.43, Z68.44, Z68.45
<p>Sexually transmitted diseases Intensive behavioral counseling for sexually active teenagers and adults at high risk of contracting sexually transmitted diseases.</p>	99401 99402 99403 99404	Z70.0, Z70.1, Z70.2, Z70.3, Z70.8, Z70.9, Z71.7, Z72.51, Z72.52, Z72.53

Preventive Services for Adults	Procedure Codes and Facility Codes	Diagnosis Codes
<p>Statins to prevent cardiovascular events in adults: Preventive Medications The USPSTF recommends prescribing a statin for the primary prevention of cardiovascular disease (CVD) for adults aged 40 to 75 years who have one (1) or more CVD risk factors (for example, dyslipidemia, diabetes, hypertension, or smoking) and an estimated 10-year cardiovascular event risk of 10% or greater.</p>	Procedure 80061 82465 83718 Facility 0301	Z13.220
<p>Syphilis screening tests in teenagers and adults who are not currently pregnant The USPSTF recommends screening for syphilis infection in asymptomatic and nonpregnant teenagers and adults who are at increased risk for infection.</p>	Procedure 86592 86780 Facility 0302	Z11.3, Z72.51, Z72.52, Z72.53
<p>Tobacco use and drug use cessation: Adults who are not currently pregnant The USPSTF recommends that physicians ask all adults about their tobacco use, discourage this habit, and offer behavioral interventions and smoking cessation drugs approved by the Food and Drug Administration (FDA). For those using products to cease tobacco use, this plan covers the supply of FDA-approved smoking cessation medications for ninety (90) consecutive days in a single attempt and for up to two (2) attempts per year.</p>	99406 99407	F17.200, F17.201, F17.203, F17.208, F17.209, F17.210, F17.211, F17.213, F17.218, F17.219, F17.220, F17.221, F17.223, F17.228, F17.229, F17.290, F17.291, F17.293, F17.298, F17.299, Z87.891
<p>TB Screening Test: adults Screening for tuberculosis infection in high-risk populations.</p>	86481	Z11.7
<p>Harmful alcohol use: adults Screening for harmful alcohol use at primary care facilities in adults over 18 years old, including pregnant members, by providing brief behavioral counseling interventions to reduce harmful alcohol consumption for people who engage in dangerous or risky consumption.</p>	99408 99409	F10.10, F10.120, F10.121, F10.129, F10.14, F10.150, F10.151, F10.159, F10.180, F10.181, F10.182, F10.188, F10.19, F11.10, F12.10, F13.10, F13.90, F15.90, F16.90, F17.290, Z71.41, Z71.51, Z71.6, Z87.891, Z91.89
<p>Harmful drug use in adults Screening for harmful drug use in adults 18 years of age or older by providing brief behavioral counseling interventions. The screening should be performed when services for accurate diagnosis, effective treatment, and appropriate care can be offered or referred.</p>	99408 99409	F11.10, F12.10, F13.10, F13.90, F15.90, F16.90, F17.290, Z71.41, Z71.51, Z71.6, Z87.891, Z91.89

Preventive Services for Adults, including Pregnancies

Preventive Services for Adults, including Pregnancies	Procedure Codes and Facility Codes	Diagnosis Codes
<p>Screening for bacteria in pregnant individuals The USPSTF recommends performing urine culture screening for asymptomatic bacteriuria in pregnant individuals.</p>	<p>87086 87088 87081 87084</p>	<p>O09.00, O09.01, O09.02, O09.03, O09.10, O09.11, O09.12, O09.13, O09.211, O09.212, O09.213, O09.219, O09.291, O09.292, O09.293, O09.299, O09.30, O09.31, O09.32, O09.33, O09.40, O09.41, O09.42, O09.43, O09.511, O09.512, O09.513, O09.519, O09.521, O09.522, O09.523, O09.529, O09.611, O09.612, O09.613, O09.619, O09.621, O09.622, O09.623, O09.629, O09.811, O09.812, O09.813, O09.819, O09.821, O09.822, O09.823, O09.829, O09.90, O09.91, O09.92, O09.93, Z33.1, Z34.00, Z34.01, Z34.02, Z34.03, Z34.80, Z34.81, Z34.82, Z34.83, Z34.90, Z34.91, Z34.92, Z34.93</p>
<p>BRCA: Risk Assessment Primary care providers must screen people who have had relatives with breast, ovarian, fallopian, or peritoneal cancer, using tools designed to identify family history that could be associated with an increased risk for potentially harmful mutations in the breast cancer susceptibility genes (BRCA1 or BRCA2). Members whose tests suggest they might be at risk must receive genetic counseling and, if prescribed as a result, the BRCA test.</p>	<p>99385 99386 99387 99395 99396 99397 81212 81215 81216 81217 81162 81163 81164 81165 81166 96040 99401 99402 99403 99404</p>	<p>Z31.5 Z80.3 Z80.41</p>
<p>Breast cancer: preventive drugs to reduce risk The USPSTF recommends that physicians offer to prescribe risk-reducing medications, such as tamoxifen, raloxifene, or aromatase inhibitors, to women who are at increased risk for breast cancer who are 35 years of age or older and at low risk of having adverse reactions to the medications.</p>	<p>NDC Tamoxifen, Raloxifen</p>	<p>Z15.01 Z80.3</p>

Preventive Services for Adults, including Pregnancies	Procedure Codes and Facility Codes	Diagnosis Codes
<p>Breast Cancer Screening Pursuant to Act No. 10 of January 3, 2020, “Law on the Right to Effective Breast Cancer Screening,” establishes that the following must be included among the preventive care benefits:</p> <ul style="list-style-type: none"> • One mammography, for members who are thirty-five (35) to thirty-nine (39) years old. • One (1) annual mammography for members who are forty (40) years or age or older. • One (1) annual mammography, follow-up treatment, or supplementary diagnostic tests (MRI/sonomammograms), for members who are forty (40) years of age or older whose breast tissue is considered to be heterogeneously dense or extremely dense. • One (1) annual mammography, follow-up treatment, or supplementary diagnostic tests, for women who are at high risk of developing breast cancer due to their family history, their own history as a cancer patient, the presence of high-risk markers in their genetic profile, or any other factor determined by their physician. 	<p>Procedure: 77067</p> <p>Facility: 0329</p>	<p>Z12.31</p>
<p>Breastfeeding Comprehensive breastfeeding support services (including consultation, counseling, education by physicians and peer support services, and breastfeeding equipment and supplies) during the prenatal, perinatal, and postpartum periods to optimize successful breastfeeding initiation and maintenance. Breastfeeding equipment and supplies include, but are not limited to, double electric breast pumps (including pump parts and maintenance) and breast milk storage supplies. Access to double electric breast pumps must be prioritized in order to optimize breastfeeding and will not be based on a previous problem with manual breast pumps. The breastfeeding team may also include equipment and supplies as clinically indicated to support dyads with breastfeeding difficulties and those in need of additional services. No monetary limits apply.</p> <p>Breastfeeding equipment and supplies cannot be provided through reimbursement or optional major medical coverage.</p>	<p>99401 99402 99403 99404 E0603 (Mod UN, RR, UE) E0602 (Mod UN, RR, UE)</p>	<p>Z39.1</p>
<p>Cervical cancer screening Cervical cancer screening for members who are 21 to 29 years old, with the Papanicolaou test every three (3) years. For members from 30 to 65 years old, one test every three (3) years with cervical cytology, every five (5) years with the high-risk human papillomavirus (hrHPV) test, or every five (5) years with the hrHPV test in combination with cytology.</p>	<p>G0101 G0123 G0124 G0141 G0143 G0144 G0145 G0147 G0148</p> <p>G0476</p> <p>87624 87625</p>	<p>Z01.411 Z01.419 Z12.4 Z12.72</p> <p>Z01.411, Z01.419 Z12.4</p> <p>Z00.00, Z00.01, Z01.411, Z01.419, Z11.51, Z12.4</p>
<p>Chlamydia screening Screening for chlamydia for all members who are pregnant and for members who are sexually active and under 24 years old, or older if at a high risk for infection.</p>	<p>Procedure 87490 87491</p> <p>Facility 0306</p>	<p>Z72.51 Z72.52 Z72.53 Z11.3</p>

Preventive Services for Adults, including Pregnancies	Procedure Codes and Facility Codes	Diagnosis Codes
<p>Contraceptive Methods FDA-approved contraceptive methods, sterilization procedures, screening, counseling, and education for all members of reproductive age, as prescribed. The insertion and removal of any device are covered. Follow-up visits are also considered part of contraceptive care.</p>	A4261 A4264 A4266 A4268 A4269 J7300 J7296 J7298 J7304 J7306 J7307 S4981 S4989 S4993 11980 J7294 11980 11981 11982 11983 57170 58300 58301 58565 58600 58605 58611 58615 58661 58670 58671 58720 96372 55250	Z30.011, Z30.012, Z30.013, Z30.014, Z30.018, Z30.019, Z30.02, Z30.09, Z30.2, Z30.40, Z30.41, Z30.42, Z30.430, Z30.431, Z30.432 Z30.433, Z30.49, Z30.8 Z30.432 APPLY ONLY FOR CPT 58301
<p>Healthy weight and weight gain counseling for pregnant individuals Behavioral counseling interventions for teenagers and adults to promote a healthy increase in weight and prevent excessive weight gain during pregnancy.</p>	99401 99402 99403 99404	Z34.81 Z34.82 Z34.83
<p>Intimate partner violence, elder abuse, and abuse of vulnerable adults: Screening: women of reproductive age Screening for violence, such as domestic violence, in the intimate relationships of members of reproductive age, and offering and referring individuals who tested positive to the screening to intervention services.</p>	99401 99402 99403 99404	Z00.00, Z00.01, Z04.9
<p>Folic acid supplementation to prevent neural tube defects: preventive medicine For all people who are planning or who may become pregnant, they must take a daily folic acid supplement of 0.4 to 0.8 mg (400 to 800 ug).</p>	N/A	N/A
<p>Gestational Diabetes Mellitus The Women's Preventive Services Initiative (WPSI) recommends screening pregnant women for gestational diabetes mellitus (GDM) after 24 weeks of gestation (preferably between 24 and 28 weeks of gestation) to prevent adverse birth outcomes. The WPSI suggests that pregnant women with risk factors for type 2 diabetes or GDM be screened before 24 weeks of gestation —ideally at the first prenatal visit.</p>	82947 82950 83036	O09.00, O09.01, O09.02, O09.03, O09.10, O09.11, O09.12, O09.13, O09.291, O09.292, O09.293, O09.299, O09.40, O09.41, O09.42, O09.43, Z13.1, Z33.1, Z34.00, Z34.01, Z34.02, Z34.03, Z34.80, Z34.81, Z34.82, Z34.83, Z34.90, Z34.91, Z34.92, Z34.93
<p>Gonorrhea screening Screening for gonorrhea in individuals who are sexually active and no more than 24 years old, or older if at a high risk for infection.</p>	Procedure 87590 87591	O09.00, O09.01, O09.02, O09.03, O09.10, O09.11, O09.12, O09.13, O09.211, O09.212,

Preventive Services for Adults, including Pregnancies	Procedure Codes and Facility Codes	Diagnosis Codes
	Facility 0306	O09.213, O09.219, O09.291, O09.292, O09.293, O09.299, O09.30, O09.31, O09.32, O09.33, O09.40, O09.41, O09.42, O09.43, O09.511, O09.512, O09.513, O09.519, O09.521, O09.522, O09.523, O09.529, O09.611, O09.612, O09.613, O09.619, O09.621, O09.622, O09.623, O09.629, O09.811, O09.812, O09.813, O09.819, O09.821, O09.822, O09.823, O09.829, O09.90, O09.91, O09.92, O09.93, Z11.3, Z33.1, Z34.00, Z34.01, Z34.02, Z34.03, Z34.80, Z34.81, Z34.82, Z34.83, Z34.90, Z34.91, Z34.92, Z34.93, Z72.51, Z72.52, Z72.53
Hepatitis B screening: pregnant members For pregnant members, screening for Hepatitis B virus infection during the first prenatal visit.	Procedure 87340 Facility 0306	O09.00, O09.01, O09.02, O09.03, O09.10, O09.11, O09.12, O09.13, O09.211, O09.212, O09.213, O09.219, O09.291, O09.292, O09.293, O09.299, O09.30, O09.31, O09.32, O09.33, O09.40, O09.41, O09.42, O09.43, O09.511, O09.512, O09.513, O09.519, O09.521, O09.522, O09.523, O09.529, O09.611, O09.612, O09.613, O09.619, O09.621, O09.622, O09.623, O09.629, O09.811, O09.812, O09.813, O09.819, O09.821, O09.822, O09.823, O09.829, O09.90, O09.91, O09.92, O09.93, Z33.1, Z34.00, Z34.01, Z34.02, Z34.03, Z34.80, Z34.81, Z34.82, Z34.83, Z34.90, Z34.91, Z34.92, Z34.93, Z11.59

Preventive Services for Adults, including Pregnancies	Procedure Codes and Facility Codes	Diagnosis Codes
<p>Screening test for human immunodeficiency virus (HIV): Pregnant members</p> <p>Physicians must screen all pregnant members for HIV, including those who come in for delivery and have not been tested and whose HIV status is unknown. The following tests are covered at no copayment for pregnant individuals:</p> <ol style="list-style-type: none"> First HIV test during the first trimester of gestation or on the first prenatal visit, and Second test during the third trimester of gestation (between 28 and 34 weeks of gestation). 	<p>Procedure 86701 86702 86703 86689 87390</p> <p>Facility 0302</p>	<p>O09.00, O09.01, O09.02, O09.03, O09.10, O09.11, O09.12, O09.13, O09.211, O09.212, O09.213, O09.219, O09.291, O09.292, O09.293, O09.299, O09.30, O09.31, O09.32, O09.33, O09.40, O09.41, O09.42, O09.43, O09.511, O09.512, O09.513, O09.519, O09.521, O09.522, O09.523, O09.529, O09.611, O09.612, O09.613, O09.619, O09.621, O09.622, O09.623, O09.629, O09.811, O09.812, O09.813, O09.819, O09.821, O09.822, O09.823, O09.829, O09.90, O09.91, O09.92, O09.93, Z33.1, Z34.00, Z34.01, Z34.02, Z34.03, Z34.80, Z34.81, Z34.82, Z34.83, Z34.90, Z34.91, Z34.92, Z34.93</p>
<p>Human Immunodeficiency Virus (HIV) screening test</p> <ul style="list-style-type: none"> All adolescents and adult women (15 years and older) should be tested for HIV at least once in their lifetime. Earlier or additional screening tests should be based on risk, and annual or more frequent retesting may be appropriate starting at age 13 for adolescents and adult women at increased risk of HIV infection. Risk assessment and HIV prevention education beginning at age 13 and continuing as determined by risk. An HIV screening test is recommended for all pregnant women at the start of prenatal care, with retesting during the pregnancy based on risk factors. Rapid HIV testing is recommended for pregnant women who have gone into active labor and whose HIV status is unknown. Screening during pregnancy can help prevent vertical transmission. 	<p>86701 86702 86703 86689 87390 99401 99402 99403 99404</p>	<p>Z70.0, Z70.1, Z70.2, Z70.3, Z70.8, Z70.9, Z71.7, Z72.51, Z72.52, Z72.53</p>
<p>Prevention of obesity in middle-aged women.</p> <p>Counseling middle-aged women (40 to 60 years old) who have a normal or overweight body mass index (BMI) (18.5 – 29.9 kg/m²) to help them maintain their weight or limit their weight gain in order to prevent obesity. These counseling services may include a one-on-one discussion on healthy nutrition and physical activity.</p>	<p>99386</p>	<p>Z00.00, Z00.01, Z01.411, Z01.419, Z04.9, Z12.11, Z12.12, Z12.13, Z12.31, Z12.6, Z68.20, Z68.21, Z68.22, Z68.23, Z68.25, Z68.26, Z68.27, Z68.28, Z68.29</p>

Preventive Services for Adults, including Pregnancies	Procedure Codes and Facility Codes	Diagnosis Codes
Counseling and Screening for Human Immunodeficiency Virus Annual counseling and screening for human immunodeficiency virus infection for all sexually active women.	86701 86702 86703 86689 87390 99401 99402 99403 99404	Z70.0, Z70.1, Z70.2, Z70.3, Z70.8, Z70.9, Z71.7, Z72.51, Z72.52, Z72.53
Osteoporosis screening: postmenopausal members under 65 years old at a higher risk for osteoporosis Screening for osteoporosis with a bone densitometry test to prevent osteoporosis fractures in postmenopausal members under 65 years old who are at a higher risk for osteoporosis, as determined through a formal clinical risk assessment tool.	Procedure 77080 77081 Facility 0320	Z13.820
Osteoporosis screening: members over 65 years old Screening for osteoporosis with a bone densitometry test to prevent osteoporosis fractures in members over 65 years old.	Procedure 77080 77081 Facility 0320	Z13.820
Perinatal Depression: counseling and intervention Clinical staff are advised to provide interventional counseling or refer pregnant or postpartum members who are at a risk for perinatal depression.	96160 96161 99384 99385 99386 99387 99394 99395 99396 99397	Z13.32
Preeclampsia prevention: aspirin Use of low-dose aspirin (81mg/d) as preventive medication after the 12 th week of gestation in members at high risk for preeclampsia.	N/A	O14.10, O14.12 y O14.13
Hypertensive disorders of pregnancy: asymptomatic pregnant individuals Screening for preeclampsia in people using blood pressure monitoring throughout the entire pregnancy.	N/A	O14.10, O14.12 y O14.13
Rh(D) Incompatibility Screening Rh(D) blood type and antibody tests for all pregnant members during their pregnancy at the first prenatal visit. Includes repeating the antibody test for pregnant members who have Rh-negative blood but aren't Rh-sensitized, sometime between 24 and 28 weeks of gestation, unless the biological father is also Rh-negative.	Procedure 86901 Facility 0301	O09.00, O09.01, O09.02, O09.03, O09.10, O09.11, O09.12, O09.13, O09.211, O09.212, O09.213, O09.219, O09.291, O09.292, O09.293, O09.299, O09.30, O09.31, O09.32, O09.33, O09.40, O09.41, O09.42, O09.43, O09.511, O09.512, O09.513, O09.519, O09.521, O09.522, O09.523, O09.529, O09.611, O09.612, O09.613, O09.619, O09.621, O09.622,

Preventive Services for Adults, including Pregnancies	Procedure Codes and Facility Codes	Diagnosis Codes
		O09.623, O09.629, O09.811, O09.812, O09.813, O09.819, O09.821, O09.822, O09.823, O09.829, O09.90, O09.91, O09.92, O09.93, Z33.1, Z34.00, Z34.01, Z34.02, Z34.03, Z34.80, Z34.81, Z34.82, Z34.83, Z34.90, Z34.91, Z34.92, Z34.93
<p>Anxiety screening tests The Women's Preventive Services Initiative recommends anxiety screening tests for teenage and adult women, including those who are pregnant or postpartum. The optimal intervals for screening are unknown, and clinical judgment should be used to determine the frequency for these assessments. Given the high prevalence of anxiety disorders, the lack of recognition in clinical practice, and the multiple problems associated with untreated anxiety, physicians should consider screening women who have not been recently screened.</p>	99385 99386	Z13.39
<p>Screening for diabetes mellitus after pregnancy The Women's Preventive Services Initiative (WPSI) recommends screening for type 2 diabetes among women with a history of gestational diabetes mellitus (GDM) who are not currently pregnant and have not been previously diagnosed with type 2 diabetes.</p> <p>Ideally, initial tests must be performed within the first year postpartum and may be performed as soon as 4 to 6 weeks after delivery.</p> <p>For women who were not screened in the first year postpartum or whose initial result was negative, the postpartum screening test should be performed at least every 3 years for a minimum of 10 years after pregnancy. For those who test positive for diabetes in the early postpartum period, the test should be repeated at least 6 months postpartum to confirm the diabetes diagnosis regardless of the initial test type (for example, fasting plasma glucose, hemoglobin A1C, glucose tolerance test). Repeat testing is also indicated for women tested with hemoglobin A1C in the first 6 months postpartum, regardless of whether the test results were positive or negative, because hemoglobin A1C testing is less accurate during the first 6 months postpartum.</p> <p>For women who were not screened in the first year postpartum or whose initial result was negative, the postpartum screening test should be performed at least every 3 years for a minimum of 10 years after pregnancy. For those who test positive for diabetes in the early postpartum period, the test should be repeated at least 6 months postpartum to confirm the diabetes diagnosis regardless of the initial test type (for example, fasting plasma glucose, hemoglobin A1C, glucose tolerance test).</p> <p>Repeat testing is also indicated for women tested with hemoglobin A1C in the first 6 months postpartum, regardless of whether the test results were positive or negative, because hemoglobin A1C testing is less accurate during the first 6 months postpartum.</p>	82947 82951 83037	Z13.1 Z86.32

Preventive Services for Adults, including Pregnancies	Procedure Codes and Facility Codes	Diagnosis Codes
<p>Syphilis screening during pregnancy Screening for syphilis in all members during pregnancy.</p>	<p>Procedure 86592 86780</p> <p>Facility 0302</p>	<p>O09.00, O09.01, O09.02, O09.03, O09.10, O09.11, O09.12, O09.13, O09.211, O09.212, O09.213, O09.219, O09.291, O09.292, O09.293, O09.299, O09.30, O09.31, O09.32, O09.33, O09.40, O09.41, O09.42, O09.43, O09.511, O09.512, O09.513, O09.519, O09.521, O09.522, O09.523, O09.529, O09.611, O09.612, O09.613, O09.619, O09.621, O09.622, O09.623, O09.629, O09.811, O09.812, O09.813, O09.819, O09.821, O09.822, O09.823, O09.829, O09.90, O09.91, O09.92, O09.93, Z11.3, Z33.1, Z34.00, Z34.01, Z34.02, Z34.03, Z34.80, Z34.81, Z34.82, Z34.83, Z34.90, Z34.91, Z34.92, Z34.93, Z72.51, Z72.52, Z72.53</p>
<p>Urinary incontinence screening tests in women The Women's Preventive Services Initiative recommends screening women to detect urinary incontinence as a preventive service. Factors associated with a higher risk for urinary incontinence include a higher parity, advancing age, and obesity; however, these factors should not be used to limit screening.</p> <p>Several screening tools have demonstrated moderate to high accuracy in identifying urinary incontinence in women. Although the minimum screening intervals are unknown, given the prevalence of urinary incontinence, it is advisable to perform the test annually since many women do not exhibit symptoms, and the multiple risk factors associated with incontinence change frequently.</p>	<p>99384 99385 99386 99387 99394 99395 99396 99397</p>	<p>Z01.419</p>
<p>Tobacco use and smoking cessation for pregnant members Physicians must ask all pregnant members about their tobacco use, advise them to stop, and provide behavioral interventions to help smoking members cease their tobacco consumption.</p>	<p>99406 99407</p>	<p>F17.200, F17.201, F17.203, F17.208, F17.209, F17.210, F17.211, F17.213, F17.218, F17.219, F17.220, F17.221, F17.223, F17.228, F17.229, F17.290, F17.291, F17.293, F17.298, F17.299, O09.511, O09.512, O09.513, O09.519, O09.521, O09.522,</p>

Preventive Services for Adults, including Pregnancies	Procedure Codes and Facility Codes	Diagnosis Codes
		O09.523, O09.529, O09.611, O09.612, O09.613, O09.619, O09.621, O09.622, O09.623, O09.629, O09.811, O09.812, O09.813, O09.819, O09.821, O09.822, O09.823, O09.829, O99.330, O99.331, O99.332, O99.333, O99.334, O99.335, Z34.00, Z34.01, Z34.02, Z34.03, Z34.80, Z34.81, Z34.82, Z34.83, Z34.90, Z34.91, Z34.92, Z34.93, Z87.891
Preventive visits for members Annual preventive visit (depending on the member's health needs and other risk factors) so that adult members can access the recommended preventive services adequate for their age, including essential prenatal care and services. Whenever appropriate, this annual preventive visit must include other listed preventive services. Should the physician determine that the patient requires additional visits, these will be covered without copayment.	99384 99385 99386 99387 99394 99395 99396 99397	Z00.00, Z00.01, Z01.411, Z01.419, Z04.9, Z12.11, Z12.12, Z12.13, Z12.31, Z12.6

Preventive Services for Children

A preventive health care visit for children normally includes the following services: medical history, measurements, sensory screening, developmental/behavioral assessment, physical examination, anticipatory guidance (such as nutritional counseling), and dental referrals, among others. The following services are available to the minor, based on age and other established guidelines as indicated below:

Preventive Services for children	Procedure Codes and Facility Codes	Diagnosis Codes
Anemia / Iron Deficiency Perform risk assessment or screening as appropriate, as recommended in the current edition of the AAP Pediatric Nutrition: Policy of the American Academy of Pediatrics (Iron Supplementation). Iron supplement for children from 4 months to 21 years of age who are at risk for anemia.	99381 99382 99383 99384 99385 99391 99392 99393 99394 99395	Z00.110 Z00.111 Z00.121 Z00.129,
Anxiety in children and teenagers: Screening: children and teenagers between 8 and 18 years old The USPSTF recommends screening for anxiety in children and teenagers from 8 to 18 years old.	99383 99384 99385	Z13.39
Behavioral, social, and emotional screening The American Academy of Pediatrics (AAP) recommends performing an annual assessment from birth to 21 years of age.	99381 99382 99383 99384 99385	Z65.8

Preventive Services for children	Procedure Codes and Facility Codes	Diagnosis Codes
Bilirubin screening Screening for newborns.	99461	Z00.110, Z00.111
Blood pressure Screening for minors: 0-11 months, 1-4 years, 5-10 years, 11-14 years, 15-17 years.	99381 99382 99383 99384 99385 99391 99392 99393 99394	Z00.110 Z00.111 Z00.121 Z00.129
Blood test Screening for newborns.	99461	Z00.110, Z00.111
Cervical displacement Screening for sexually active members.	N/A	
Screening for depression and suicide risk in children and teenagers The American Academy of Pediatrics (AAP) and the USPSTF recommend screening for major depressive disorder (MDD) in teenagers from 12 to 21 years old.	99384 99385 99394 96160 96161	Z13.89
Developmental screening and monitoring Screening for children under 3 years of age and monitoring throughout childhood.	99382 99383 99392 99393	Z00.121 Z00.129
Dyslipidemia Screening for minors, once between 9 to 11 years of age, and once again between 17 to 21 years of age. Screening for minors at risk for lipid disorders. Ages: 1-4 years, 5-10 years, 11-14 years, 15-17 years.	This recommendation has a Grade (I) statement. USPTF concludes that the current evidence is insufficient to assess the balance of benefits and harms of the service.	
Eye prophylaxis for gonorrhea: preventive medication Topical eye medication to prevent gonorrhea in newborns.	N/A	N/A
Hearing Hearing screening for all newborns and for minors, once between 11 to 14 years old, once between 15 to 17 years old, and once between 18 to 21 years old.	92551	Z00.00, Z00.01, Z00.121, Z00.129, Z01.10, Z01.118
Growth in terms of height, weight, and body mass index Screening for minors. Ages: 0-11 months, 1-4 years, 5-10 years, 11-14 years, 15-17 years.	99382 99383 99392 99393	Z00.121 Z00.129
Hematocrit or hemoglobin screening Screening for all minors when there is a risk factor.	99382 99383 99392 99393	Z00.121 Z00.129

Preventive Services for children	Procedure Codes and Facility Codes	Diagnosis Codes
<p>Sickle cell disease (hemoglobinopathy) Screening for sickle cell disease in newborns.</p>	<p>Procedure 83020</p> <p>Facility 0306, 0301</p>	<p>O09.00, O09.01, O09.02, O09.03, O09.10, O09.11, O09.12, O09.13, O09.211 O09.212, O09.213, O09.219, O09.291, O09.292, O09.293, O09.299, O09.30, O09.31 O09.32, O09.33, O09.40, O09.41, O09.42, O09.43, O09.511, O09.512, O09.513, O09.519, O09.521, O09.522, O09.523, O09.529, O09.611, O09.612, O09.613, O09.619, O09.621, O09.622, O09.623, O09.629, O09.811, O09.812, O09.813, O09.819, O09.821, O09.822, O09.823, O09.829, O09.90, O09.91, O09.92, O09.93, Z33.1, Z34.00, Z34.01, Z34.02, Z34.03, Z34.80, Z34.81, Z34.82, Z34.83, Z34.90, Z34.91, Z34.92, Z34.93</p>
<p>Hepatitis B screening The USPSTF and the American Academy of Pediatrics (AAP) recommend screening for hepatitis B virus (HBV) infection in newborns up to early adulthood (21 years old) who are at high risk for infection.</p>	<p>Procedure 87340</p> <p>Facility 0306</p>	<p>O09.00, O09.01, O09.02, O09.03, O09.10, O09.11, O09.12, O09.13, O09.211, O09.212, O09.213, O09.219, O09.291, O09.292, O09.293, O09.299, O09.30, O09.31, O09.32, O09.33, O09.40, O09.41, O09.42, O09.43, O09.511, O09.512, O09.513, O09.519, O09.521, O09.522, O09.523, O09.529, O09.611, O09.612, O09.613, O09.619, O09.621, O09.622, O09.623, O09.629, O09.811, O09.812, O09.813, O09.819, O09.821, O09.822, O09.823, O09.829, O09.90, O09.91, O09.92, O09.93, Z33.1, Z34.00, Z34.01, Z34.02, Z34.03, Z34.80, Z34.81, Z34.82, Z34.83, Z34.90, Z34.91, Z34.92, Z34.93, Z11.59</p>

Preventive Services for children	Procedure Codes and Facility Codes	Diagnosis Codes
	90682	Z23
	90685	Z23
	90686	Z23
	90687	Z23
	90688	Z23
	90696	Z23
	90698	Z23
	90700	Z23
	90702	Z23
	90707	Z23
	90713	Z23
	90714	Z23
	90715	Z23
	90716	Z23
	90723	Z23
	90732	Z23
	90733	Z23
	90734	Z23
	90740	Z23
	90743	Z23
	90744	Z23
	90746	Z23
	90748	Z23
	90671	Z23
Lead Screening for minors from 1 to 6 years old with high levels of lead in their blood who are at a moderate-to-high risk, and for pregnant members exhibiting no symptoms.	83655	Z13.88 Z72.89
Postpartum depression Screening for mothers of newborns during their visits at 1, 2, 4 and 6 months.	96160 96161 99384 99385 99386 99387 99394 99395 99396 99397	Z13.32
Medical history For any minor during development, from 0 to 21 years old.	99381 99382 99383 99384 99385	Z00.121 Z00.129
Screening for obesity in minors and teenagers For minors and teenagers (6 years old and up), intensive comprehensive behavioral interventions to promote an improvement in the child's weight.	99383 99384 99393 99394	E66.01, Z68.53, Z68.54
Oral health Risk assessment for minors from 0 to 11 months old, 1 to 4 years old, and 5 to 10 years old.	99381 99391 99382 99392	Z29.3

Preventive Services for children	Procedure Codes and Facility Codes	Diagnosis Codes
<p>Phenylketonuria (PKU) screening for newborns Screening for phenylketonuria (PKU) in newborns.</p>	<p>Procedure 84030</p> <p>Facility 301</p>	<p>Z13.228</p>
<p>Prevention of tooth decay in children under 5 years of age: Screening and interventions: children under 5 years of age The USPSTF recommends that primary care clinicians prescribe oral fluoride supplementation starting at age 6 months for children whose water supply is deficient in fluoride. The American Academy of Pediatrics (AAP) also recommends considering adding an oral fluoride supplement if the main water source is fluoride-deficient. The USPSTF recommends that primary care clinicians apply fluoride varnish to the primary teeth of all infants and children starting at the age of primary tooth eruption. Once the tooth is in place, apply fluoride varnish to all children between 3 to 6 months of age via primary care or at the dental office, based on risk for cavities.</p>	<p>99188 99381 99391 99382 99392</p>	<p>Z29.3</p>
<p>Skin cancer: Counseling Counseling for minors, teenagers, and young adults with white skin, aged 6 months to 24 years old, to minimize their exposure to ultraviolet radiation and reduce their risk for skin cancer.</p>	<p>D1206</p>	<p>N/A</p>
<p>Sudden cardiac arrest and sudden cardiac death The American Academy of Pediatrics (AAP) recommends counseling on the risks of sudden cardiac arrest and sudden cardiac death, and this has been added for those 11 to 21 years of age (to account for the range in which risk counseling may take place) to be consistent with AAP's policy ("Sudden Death in the Young: Information for the Primary Care Provider"). Perform an assessment, as appropriate.</p>	<p>99401 99402</p>	<p>Z71.89</p>
<p>Tobacco use in minors and teenagers Interventions, including education and counseling, for minors and teenagers, to prevent the start of tobacco use.</p>	<p>99406 99407</p>	<p>F17.200, F17.201, F17.203, F17.208, F17.209, F17.210, F17.211, F17.213, F17.218, F17.219, F17.220, F17.221, F17.223, F17.228, F17.229, F17.290, F17.291, F17.293, F17.298, F17.299, O09.511, O09.512, O09.513, O09.519, O09.521, O09.522, O09.523, O09.529, O09.611, O09.612, O09.613, O09.619, O09.621, O09.622, O09.623, O09.629, O09.811, O09.812, O09.813, O09.819, O09.821, O09.822, O09.823, O09.829, O99.330, O99.331, O99.332, O99.333, O99.334, O99.335, Z34.00, Z34.01, Z34.02,</p>

Preventive Services for children	Procedure Codes and Facility Codes	Diagnosis Codes
		Z34.03, Z34.80, Z34.81, Z34.82, Z34.83, Z34.90, Z34.91, Z34.92, Z34.93, Z87.891
Tobacco, alcohol, and drug use Screening for minors aged 11 to 21 years old.	99408 99409	F11.10, F12.10, F13.10, F13.90, F15.90, F16.90, F17.290, Z71.41, Z71.51, Z71.6, Z87.891, Z91.89
Tuberculosis Test for minors at high risk for tuberculosis. Ages: 0-11 months, 1-4 years, 5-10 years, 11-14 years, 15-21 years.	86580	Z11.1
Vision screening: minors Vision screening test, at least once (1) for all minors between 3 and 5 years old to detect amblyopia or risk factors.	99173	Z01.00, Z01.01, Z13.5, Z00.121, Z00.129

Triple-S Salud, Inc. cumple con las leyes federales aplicables de derechos civiles y no discrimina en base a raza, color, origen de nacionalidad, edad, discapacidad, o sexo. *Triple-S Salud, Inc. complies with applicable federal civil rights laws and does not discriminate because of race, color, national origin, age, disability, or sex. ATTENTION: If you speak English, language assistance services, free of charge, are available to you. Call 787-774-6060, (TTY/TDD), 787-792-1370 or 1-866-215-1999. Free of charge 1-800-981-3241. If you are a federal employee or retiree call 787-774-6081, Toll Free 1-800-716-6081; (TTY / TDD) 787-792-1370; Toll-Free 1-866-215-1999* ATENCIÓN: si habla español, tiene a su disposición servicios gratuitos de asistencia lingüística llame al 787-774-6060, Libre de costo 1-800-981-3241. (TTY/TDD) al 787-792-1370 o 1-866-215-1999. Si es empleado o retirado federal llame al 787-774-6081, libre de costo 1-800-716-6081; (TTY / TDD) 787-792-1370; libre de costo 1-866-215-1999. Concesionario Independiente de BlueCross BlueShield Association. TSS-PROD- 5442-2024-A

